



TITLE:

Historical perspective on the
synonymization of the four major pest
species belonging to the *Bactrocera dorsalis*
species complex (Diptera, tephritidae)

AUTHOR(S):

Hee, Alvin K W; Wee, Suk Ling; Nishida, Ritsuo; Ono,
Hajime; Hendrichs, Jorge; Haymer, David S.; Tan,
Keng Hong

CITATION:

Hee, Alvin K W ...[et al]. Historical perspective on the synonymization of the four major
pest species belonging to the *Bactrocera dorsalis* species complex (Diptera, tephritidae).
ZooKeys 2015, 540: 323-338

ISSUE DATE:

2015-11-26

URL:

<http://hdl.handle.net/2433/226725>

RIGHT:

Copyright Alvin K.W. Hee et al. This is an open access article distributed
under the terms of the Creative Commons Attribution License (CC BY
4.0), which permits unrestricted use, distribution, and reproduction in
any medium, provided the original author and source are credited.

Historical perspective on the synonymization of the four major pest species belonging to the *Bactrocera dorsalis* species complex (Diptera, Tephritidae)

Alvin K.W. Hee¹, Suk-Ling Wee^{2,3}, Ritsuo Nishida⁴, Hajime Ono⁴,
Jorge Hendrichs⁵, David S. Haymer⁶, Keng-Hong Tan⁷

1 Department of Biology, Faculty of Science, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia **2** School of Environmental and Natural Resource Sciences, **3** Centre of Insect Systematics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia **4** Laboratory of Chemical Ecology, Kyoto University, Kyoto, Japan **5** FAO/IAEA Joint Division of Nuclear Techniques in Food and Agriculture, Vienna, Austria **6** Department of Cell and Molecular Biology, University of Hawaii, Honolulu, HI, United States of America **7** Tan Hak Heng Co., Penang, Malaysia

Corresponding author: Alvin K.W. Hee (alvinhee@upm.edu.my)

Academic editor: A. Clarke | Received 23 April 2015 | Accepted 18 August 2015 | Published 26 November 2015

<http://zoobank.org/E50B29DD-BAEF-4404-9901-5A45DA1B6837>

Citation: Hee AKW, Wee SL, Nishida R, Ono H, Hendrichs J, Haymer DS, Tan KH (2015) Historical perspective on the synonymization of the four major pest species belonging to the *Bactrocera dorsalis* species complex (Diptera, Tephritidae). In: De Meyer M, Clarke AR, Vera MT, Hendrichs J (Eds) Resolution of Cryptic Species Complexes of Tephritid Pests to Enhance SIT Application and Facilitate International Trade. ZooKeys 540: 323–338. doi: 10.3897/zookeys.540.6028

Abstract

An FAO/IAEA-sponsored coordinated research project on integrative taxonomy, involving close to 50 researchers from at least 20 countries, culminated in a significant breakthrough in the recognition that four major pest species, *Bactrocera dorsalis*, *B. philippinensis*, *B. papayae* and *B. invadens*, belong to the same biological species, *B. dorsalis*. The successful conclusion of this initiative is expected to significantly facilitate global agricultural trade, primarily through the lifting of quarantine restrictions that have long affected many countries, especially those in regions such as Asia and Africa that have large potential for fresh fruit and vegetable commodity exports. This work stems from two taxonomic studies: a revision in 1994 that significantly increased the number of described species in the *Bactrocera dorsalis* species complex; and the description in 2005 of *B. invadens*, then newly incursive in Africa. While taxonomically valid species, many biologists considered that these were different names for one biological species. Many disagreements confounded attempts to develop a solution for resolving this taxonomic issue, before the FAO/IAEA project commenced. Crucial to understanding the success of that initiative is an accounting of the historical events and perspectives leading up to the international, multidisciplinary collaborative efforts that successfully achieved the final synonymization. This review highlights the 21 year journey taken to achieve this outcome.

Keywords

Bactrocera dorsalis species complex, *B. philippinensis*, *B. papayae*, *B. invadens*, synonymization, integrative taxonomy

Introduction

The genus *Bactrocera* Macquart of true fruit flies belongs to the Dacinae - a subfamily of Tephritidae (Drew 1989). Over 500 species have been described as belonging to this genus, making it one of the largest genera within the Tephritidae (Drew 1989). It has been estimated that total damage caused by tephritid fruit flies affecting production, harvesting, packing, and marketing of fruits globally amounts to over US\$2 billion annually (Shelly et al. 2014).

Within *Bactrocera*, the *B. dorsalis* species complex contains almost 100 species that are morphologically similar and occur in the tropics and subtropics where fruit growing areas are extensive (Drew and Romig 2013). A number of species in this complex are of economic importance and highly invasive, the best known of which is the Oriental fruit fly, *Bactrocera dorsalis* (Hendel). Closely linked in pest status with *B. dorsalis* are the now synonymized species *B. papayae* Drew & Hancock, *B. philippinensis* Drew & Hancock and *Bactrocera invadens* Drew, Tsuruta & White. *Bactrocera dorsalis* is known to cause devastating losses in fruit commodities, especially in the Asia-Pacific and Africa regions (Kawasaki et al. 1991, Ye 2001, Verghese et al. 2004, Clarke et al. 2005, De Meyer et al. 2010, Khamis et al. 2012, Li et al. 2012).

The existence of a complex of closely related, morphologically similar species to *B. dorsalis*, had been recognized for over 40 years (Hardy and Adachi 1954) before a major taxonomic revision of the group was undertaken by Drew and Hancock (1994). This revision described over 50 new species in the complex and, while the biological validity of most described species has not been questioned, this has not been the case for all taxa, particularly some of the very important pest species such as *B. papayae*. For these pest species, serious questions have been raised as to the validity over their status as separate species. This issue was accentuated when Drew et al. (2005) described *B. invadens* from Africa, a new species which again could not be consistently and readily separated from *B. dorsalis*.

The extensive similarities between *B. dorsalis* s.s and the three putative species established, *B. invadens*, *B. papayae* and *B. philippinensis*, has led to much debate on the delimitation of these species, particularly in terms of seeking congruence between the biological and taxonomic status of these entities (Tan 2003, Clarke et al. 2005, Drew et al. 2008). Numerous studies over the past two decades have contributed to our understanding of the pest species in the *B. dorsalis* complex particularly with respect to their morphological and biological attributes. Nonetheless, these studies had not reduced the complexities of the species' status, as evident from a rise in the number of

species in the *B. dorsalis* complex to almost 100 in the years since the 1994 revision in (Drew and Romig 2013).

To address and resolve the longstanding issue of species delimitation in the key pest species of the *B. dorsalis* complex, a multidisciplinary approach was adopted by an international team of more than fifty researchers from over twenty different countries. Under the auspices of the Joint FAO/IAEA Division on Nuclear Techniques in Food and Agriculture, a 5-year Coordinated Research Project (CRP) on '*Resolution of Cryptic Species Complexes of Tephritid Pests to Overcome Constraints to SIT Application and International Trade*' was established in 2010. The aim of the project was to define the species limits of pest species complexes within the Tephritidae, with the *B. dorsalis* complex identified as a priority. Studies that had been independently developed in the past, including morphometric, cytogenetic, molecular, behavioural and chemoeological datasets were re-examined, and gaps crucial for answering questions of how taxonomic species could be reconciled as biological species were filled. This project has led to the synonymization of *B. papayae*, *B. philippinensis* and *B. invadens* with *B. dorsalis*, based on the conclusion that there is insufficient evidence to maintain the former three taxa as biological species distinct from *B. dorsalis* (Schutze et al. 2015a).

It is not the aim of this paper to again provide the evidence for the synonymization of the four major pest species, as this has already been provided (Schutze et al. 2015a). Rather, it is our intent here to ensure that the long and arduous journey taken to achieve this outcome is understood. We feel that it is vitally important that the younger generation of fruit fly workers, who though armed with advanced scientific skills and techniques, will appreciate the background and good science conducted from the beginning in resolving this issue of significant transboundary importance for international agricultural trade. Thus, this paper aims to provide a historical account of the events leading to the FAO/IAEA-sponsored international efforts in resolving this prickly issue.

Taxonomic history of *B. dorsalis* complex

Before describing some of the biological insights which led to the questioning of taxonomic validity of these species, this section details the taxonomic history of the taxa of concern.

Bactrocera papayae, *B. carambolae* and *B. philippinensis*

Prior to the taxonomic revision of the *B. dorsalis* complex by Drew and Hancock (1994), the taxa endemic to the southeast Asian region of Malaysia, Indonesia and Thailand were identified as a single species, viz. *Bactrocera* (= *Dacus*) *dorsalis* (Hendel) (Hardy and Adachi 1954, Tan and Lee 1982). A second taxon was subsequently recognized, being referred to before description as Malaysian B (Drew 1991) and 'sp. near

B. dorsalis (B)' (White and Elson-Harris 1992), before being formally described as *B. papayae* Drew and Hancock (1994). This taxon was given the common name of Asian papaya fly (Drew 1997), although studies in Malaysia had shown that papaya was not the preferred host of this species, which prefers instead starfruit and banana (Tan and Nishida 1996). The detection of *B. papayae* in northern Queensland, Australia, in 1995 resulted in a successful eradication programme costing over US\$32.5 million (Fay et al. 1997, Cantrell et al. 2002).

The concern over the destructive potential of *B. papayae* also underscored the importance of another closely related species, *B. carambolae* Drew & Hancock, which itself had formerly been referred to as Malaysian A (Drew 1991) and sp. near *B. dorsalis* (A) (White and Elson-Harris 1992) and was found together with *B. papayae* in Peninsular Malaysia and southern Thailand (Clarke et al. 2001). Together, *B. papayae* and *B. carambolae* accounted for the most damage to fresh fruits in Malaysia. These species were already known to be morphologically similar and able to interbreed resulting in viable laboratory offspring with hybrid rectal pheromonal compounds, even up to the F₃ generation (Wee 2000). Natural hybrids of both *B. carambolae* and *B. papayae* possessing similar rectal pheromonal compounds to those of laboratory hybrids had also been detected from the field (Wee and Tan 2005).

Additional to 'species near *B. dorsalis* (A) and (B), was a third taxon, designated as 'sp. near *B. dorsalis* (C) (White and Elson-Harris 1992). This population was only known from the Philippines and was subsequently described by Drew and Hancock (1994) as *B. philippinensis*.

Bactrocera invadens

As for the South-east Asian pest species of the *B. dorsalis* complex, confusion also existed for *B. invadens* in Africa, a devastating pest species now widespread in Africa, which has largely displaced other long established pests such as Mediterranean fruit fly, *Ceratitidis capitata* (Wiedemann) (Hill and Treblanche 2014). Between its first detection in Kenya, 2003 and subsequent description as a new species (Drew et al. 2005), it was believed to originate from the *B. dorsalis* group, and was most likely *B. dorsalis* s.s. (Lux et al. 2003). Drew et al. (2005) and Drew et al. (2008) treated *B. invadens* as part of the *B. dorsalis* complex, but it was subsequently removed from the complex (Drew and Romig 2013).

Accumulating evidence for synonymisation – 1994 to 2010

Biological insights from chemical ecology

Efforts of fruit fly workers over two decades to resolve the biological species status of *B. papayae*, *B. philippinensis*, *B. invadens* and *B. dorsalis* started with the question from

one of us (Keng-Hong Tan) on why males of certain fly species, such as *B. dorsalis*, are so strongly attracted to methyl eugenol (ME). This research effort was partly in response to a challenge posed by N. Tanaka of the USDA Hawaiian Fruit Flies Investigations Laboratory in 1980 to figure out the role of ME in the biology of male tephritid flies. Whilst ME was first discovered a century ago as attracting male fruit flies (Howlett 1915), it was not until 60 years later that it was shown to be a highly potent attractant for *B. dorsalis* (Metcalf et al. 1975). While known to respond to ME, the question of why male flies are uniquely attracted to ME has been considered as a great mystery of tephritid biology (Cunningham 1989). ME is found as a common phenylpropanoid in numerous species of plants (>450 species in 80 families of plants) (Tan and Nishida 2012); and synthetic ME has been successfully used for monitoring and control programmes, including area-wide and border surveillance, quarantine and male annihilation programmes worldwide against many ME-responding tephritid pests of major economic importance (see reviews in Vargas et al. 2010; Shelly et al. 2014, Tan et al. 2014).

While Tan and colleagues searched for the reason as to why male *Bactrocera* respond to ME, significant insights were also gained as to the likely biological relationships between *B. dorsalis* and *B. papayae* (especially). This section describes the history of that process.

In the early 1980s work was undertaken using ME for field ecological surveys and population dynamics of *Dacus* species. This work demonstrated for the first time that temporary habituation of the male flies to ME is possible (Tan and Lee 1982; Tan 1984, 1985; Tan and Jaal 1986; Tan et al. 1987; Tan and Serit 1988). Following that and concomitant with continued ecological work involving ME, there was also interest in understanding the fate of consumed ME in relation to fruit fly physiology. Initial work by Keng-Hong Tan showed that, based on thin layer chromatography (TLC) analyses of haemolymph and digestive tissues of ME-fed and ME-deprived male flies, consumed ME was detected in the crop organ only. Furthermore, two saliva spots (with different relative mobilities from ME), that resulted from ME feeding by live sexually mature males (without prior exposure to ME), were detected on the developed-TLC plate only for extracts of ME-fed male rectal gland, crop and haemolymph (least visualized). This suggested the possibility that the rectal (pheromone) gland contained some compounds derived from the crop upon feeding of ME that were carried to the gland by the haemolymph. The transport of the ME-derived compounds to the rectal gland was subsequently confirmed (Hee and Tan 2004, 2005, 2006).

In 1986, the research collaboration of Keng-Hong Tan and Ritsuo Nishida of Kyoto University on the ecological significance of male fruit fly attractants took off after the introduction of Nishida by Professor S. Takahashi also of Kyoto University to Tan. This research collaboration resulted in the detection of the phenylpropanoids, (*E*)-coniferyl alcohol (CF) and 2-allyl-4,5-dimethoxyphenol (DMP), as sex pheromone components of male *B. dorsalis* (samples from Malaysia later known as 'Malaysian B' and then *B. papayae* - see next paragraph) following ME consumption (Nishida et al 1988a, 1988b).

During that time, *B. dorsalis* in Malaysia had been taxonomically split into two sibling species, Malaysian A and Malaysian B in 1991, and subsequently in 1994 described as *B. carambolae* and *B. papayae*, respectively. These sibling species, together with *B. dorsalis* s.s., were morphologically very similar to each other, particularly for male flies, and much work was concentrated on attempts to differentiate these species. In 1994, results from a primary chemotaxonomic diagnostic tool were presented in the 4th International Symposium of Fruit Flies of Economic Importance held in Tampa, Florida, demonstrating that CF and DMP were detected in both *B. dorsalis* (ex-Hawaii) and *B. papayae*, and with improved male mating competitiveness, following males' consumption of ME (Tan and Nishida 1996). Both the compounds were also shown to be male attractants, although CF was more attractive than DMP to conspecific females (Khoo et al. 2000). At the point source of highest concentration of these chemicals, females were observed to extrude their ovipositor, a sign of mating acceptance, confirming their role in mating behavior (Tan and Nishida 1996, Hee and Tan 1998). This information revealed the role of ME as a precursor to the ME-derived male sex pheromone components, which might be the main reason for the high mating percentages observed between the two sibling species in an outdoor cage (Tan and Nishida 1996).

The discovery that both *B. dorsalis* and *B. papayae*, having very similar morphological characters, also possessed identical pheromone components and mated readily under semi-natural conditions and producing viable offspring over generations provided the first strong evidence that the two species deserved to be considered as a single biological species. These are the basic criteria for species delimitation of Mayr's 'Biological Species Concept' (Mayr 1957).

Biochemical and genetic analyses

Following the revision of the *B. dorsalis* complex in 1994, differences in male genitalia length was used as the sole morphological basis of separating *B. papayae* from *B. dorsalis* (Drew and Hancock 1994; but see also Iwahashi 2001). Isozyme and other genetic evaluation of the two species were also conducted in an effort to provide evidence for species discrimination. However, while a discrimination was deemed possible, both species were shown to possess close genetic affinity (Yong 1994), including the lack of species-specific alleles and loci (Yong 1995). In general isozymes are now known to not be suitable markers to differentiate species due to factors such as hidden genetic variation (Behura 2006).

The development of DNA-based markers meant that a broader range of molecular tools were available to researchers in clarifying the status of species. Using ribosomal DNA markers, for example, had shown in tephritids that it was possible to differentiate distinct species within and between genera of flies, but not within a complex of closely related species (Armstrong and Cameron 2000). For example

within the *B. dorsalis* complex, no ribosomal markers in this region could be found to distinguish between *B. dorsalis*, *B. papayae* and *B. philippinensis*. In addition, another investigation that included an additional sympatric sibling species, *B. carambolae*, showed that this species also could not be discriminated by phylogenetic analyses using mitochondrial sequences (Muraji and Nakahara 2001). In this paper, the authors noted “*Although the monophylies of most of the species were supported by both topologies and bootstrap analyses, the two individuals of B. papayae did not form a clade for a species. B. papayae and its allied insects were formerly considered as a single species, B. dorsalis (Hendel)*”. The authors further stated “*Thus, B. papayae samples used in this study may be hybrids or interbreeding descendants, which might explain why the nucleotide sequences did not match the species identification based on morphological characteristics*”. In addition, Smith et al. (2003) also showed that of the four sibling species in the *B. dorsalis* complex that included *B. carambolae* and *B. caryae*, both *B. dorsalis* and *B. papayae* belonged to the same clade within the 24 *Bactrocera* species analysed. Based on a phylogenetic species theory, this suggested that *B. dorsalis* and *B. papayae* were not distinct species.

Additional molecular work on species in the *B. dorsalis* complex began with the analysis of the actin gene family in *B. dorsalis* (He and Haymer 1994). Specifically, intron sequences from these genes were used for the analysis of genetic variation in populations of *B. dorsalis* and its sibling species in the *B. dorsalis* complex (He and Haymer 1997, 2003). Introns were used primarily because, as noncoding sequences, they were typically known to be much more variable than conserved coding sequences. Because of this, it was presumed that intron sequences could serve to identify genetic markers useful for differentiating species and populations. The variable intron markers identified were also used to construct oligonucleotide arrays for rapid screening of genetic variation in populations of *B. dorsalis*, *B. papayae* and *B. carambolae* (Naeole and Haymer 2003). In this work, Naeole and Haymer showed that one of three actin alleles were identical in DNA sequence and therefore shared between *B. dorsalis* and *B. papayae*. They speculated that this result was more consistent with these taxa representing populations of the same species, rather than being distinct species. Further discussions on this possibility, as reported by Tan (2003), allowed for the beginnings of a comprehensive picture to emerge incorporating support for this idea from a wide range of datasets.

Following this, samples of *Bactrocera* flies were sent to Alfred Handler (USDA, Gainesville, Florida) from Keng-Hong Tan. The analysis of insertions of the *piggyBac* transposable element in these specimens revealed that the gene was inserted at identical loci in both *B. dorsalis* and *B. papayae*, but at different loci in other clearly distinct species (Handler et al. 2008). During the 9th Exotic Fruit Fly Symposium in Fresno, California in 2007, Handler confirmed to Tan (personal communication) that this result would be expected to occur only with extremely closely related taxa, such as between different strains of the same species that can easily interbreed and produce viable offspring, as opposed to entities representing distinct species. As a matter-of-fact, using

this same approach, Zimowska and Handler (2005) had earlier found no significant difference between that of *B. dorsalis* and *B. papayae*.

Absence of post-zygotic reproductive isolation between *B. dorsalis* and *B. papayae*

That *B. dorsalis* and *B. papayae* were not distinct species was consistent with earlier observations that interbreeding between *B. papayae*, *B. carambolae* and *B. dorsalis* results in viable offspring/hybrids (unpublished data c/f Tan 2000). Further, studies in Malaysia showed that mating compatibility between *B. dorsalis*, *B. papayae* and *B. carambolae* increased following males' consumption of ME (Wee and Tan 2000; Tan 2003). In these cases, the production of an identical phenylpropanoid, CF, functioning as a male sex pheromone component in males of those species, was suggested to be a factor in explaining the enhanced interbreeding seen in males that had consumed ME (Tan and Nishida 1996, Tan 2000). Furthermore no post-zygotic isolation was observed in matings between *B. dorsalis* (ex-Hawaii) and *B. papayae*, resulted even in F₃ hybrid offspring in the laboratory (Tan 2003).

While some authors have refuted this line of argument supporting the view that *B. dorsalis* and *B. papayae* are the same species, noting that hybridization between *Bactrocera* species is easy to achieve in laboratory cages even when using species from different subgenera (Cruickshank et al. 2001; Drew et al. 2008), interspecific matings between sibling species of *B. dorsalis*, *B. papayae* and *B. carambolae* have been reported to occur in field (McInnis et al. 1999; Wee and Tan 2000). Furthermore, in field studies, using modified ME-baited clear traps without a toxicant, the occurrence of natural hybridization between the sympatric sibling species, *B. papayae* and *B. carambolae* has been supported by captures of wild hybrids and chemotaxonomy (Wee and Tan 2005).

Persistence in maintaining the validity of *B. dorsalis* and *B. papayae* as distinct species

The failure to discover robust diagnostics markers to separate the species within the complex, and indeed the accumulating evidence that supported the idea that at least *B. dorsalis* and *B. papayae* were populations of the same species, prompted further studies by Drew et al. (2008) using additional morphometric measurements in examining the taxonomic status of a number of species in the *B. dorsalis* complex including *B. dorsalis* and *B. papayae*. In that paper, the authors argued that the accumulated evidence, including data based on morphometrics, pheromones, allozyme and nucleotide studies still supported maintaining these taxa as distinct species, even in light of the information presented in the preceding paragraphs that strongly suggested that at least *B. dorsalis* and *B. papayae* should be considered a single species. Further, whilst a review by Clarke et al. (2005) highlighted the difficulties in using morphological criteria to distinguish between those species in the *B. dorsalis* complex, particularly that of *B.*

dorsalis and *B. papayae*, the use of certain genetic diagnostic tools in the discrimination of those two species was then generally accepted. For the evidence at that time, as to why Clarke et al. and Drew et al. argued the species were valid, the reader is directed to those papers. In retrospect, and despite the strengths and weaknesses of the many individual studies and their interpretations, it seems that at the time a fully integrative taxonomic study might have contributed to resolving this issue by asking whether species that can be separated only morphologically can, in fact, be said to represent valid biological species (Clarke and Schutze 2014).

As soon as the paper by Drew et al. (2008) was published, Tan was alerted by Jorge Hendrichs of the FAO/IAEA of the authors' arguments in support of the idea that *B. dorsalis* and *B. papayae* were in fact distinct biological species. This paper included the statement that "*Although genetic similarities between allopatric populations of B. dorsalis and B. papayae have been documented (Tan, 2003), this does not cast doubt on their species status*". It is believed that the game changer to this issue was the sharp response of Keng-Hong Tan (in a form of a 4-page rebuttal letter sent via email to the journal editor) in relation to this article. In his letter, Tan pointed out that regrettably, the authors of the paper chose to ignore the increasing evidence clearly showing that the two taxa deserved recognition as one biological species. Instead, as the core focus of the article, they insisted that they were valid species based largely on morphological evaluation. The rebuttal letter was also informally sent to fruit fly researchers and members of FAO/IAEA and the International Plant Protection Convention Fruit Fly Technical Panel. Taking cognizance of the facts pointing to both *B. papayae* and *B. dorsalis* as indistinct species, we (Keng-Hong Tan, Suk-Ling Wee and Alvin Kah-Wei Hee) have already been referring to both taxa as that of *B. dorsalis* as early as 2005 in our papers (Hee and Tan 2005, 2006, Wee and Tan 2007), which were several years earlier than the implementation of the FAO/IAEA-sponsored CRP.

Furthermore, it must also be pointed out that at the same time of this fierce debate, evidence was being accumulated showing the devastation caused by another species, *B. invadens* as it invaded and spread rapidly across the continent of Africa in the early 2000s, together with the establishment of quarantine barriers by countries that did not harbour *B. invadens*, even though they harboured *B. dorsalis*.

***Bactrocera invadens* as the final impetus for coordinated international action**

The detection of an unknown pest fruit fly in Kenya, 2003, first reported as an unknown species suspected to be of Asian origin and related to the Oriental fruit fly (Lux et al. 2003), struck fear in the hearts of quarantine and trade authorities. Given much was known about the destructive losses caused by the *B. dorsalis* in many countries, the incursion of a species closely related to *B. dorsalis* was of great concern as it could potentially devastate the agricultural and fruit industries that many African countries in the tropics and subtropics depend upon. The incursion was in fact swift and destructive, with over 40 species of fruits of economic importance infested in 30 African

countries (Khamis et al. 2012). This had the international fruit fly scientific community, particularly workers in Africa and Europe, scrambling resources to seek more information on the fly's basic ecology, biology and control. Described as a new species, *B. invadens*, in 2005, over 80 refereed papers have been published on this fly since (for a brief overview, see Schutze et al. 2015b).

One major reason for the need to resolve the taxonomic status of this new pest was to confirm or reject the validity of the quarantine barriers which were established following the *B. invadens* incursion between importing countries (for example in Asia) where *B. dorsalis* was endemic and exporting African nations. In addition, an effective fruit fly management programme requires correct identification of the target pest species. In the case of *B. invadens*, while copious amounts of work have been done in understanding its ecology and biology, its true taxonomic status remained confused despite the availability of an array of modern analytical tools to aid in the identification of the species. For example, using *piggyBac* gene insertions, Zimowska and Handler (2005) found that *B. invadens* was indistinguishable from *B. dorsalis* s.s. and *B. papayae*, but this workshop abstract was not taken on to the peer reviewed literature. It must be noted that the results of Zimowska and Handler (2005) were based on the use of fly samples obtained from Tim Holler of USDA-APHIS, just prior to the formal description of these as *B. invadens*. These samples had been collected from two different populations in Tanzania (location in and around Dar Es Salaam) in December 2003 (Handler, personal communication). At the time, much of work focused on morphological evaluation, resulting in much continuing controversy on the status of this taxon as a distinct species. As an example, one of the diagnostic distinguishing characters was the appearance of varying shades of scutum colour from pale- to red-brown and almost black in *B. invadens* (Drew et al. 2005; Drew et al. 2008). The uncertainty over the species' status based on morphological features alone raised fears among some fruit fly workers that deliberation over the identity of *B. invadens* would set the stage for another grueling scientific argument, similar to that which occurred over the status of *B. dorsalis* and *B. papayae*.

The Coordinated Research Project

In conjunction with the alarming spread of *B. invadens* in Africa and doubts cast over its taxonomic status, as well the ongoing failure to resolve the species status of the Asian species, a clear need and opportunity arose for the international fruit fly research community to address the question of species boundaries in the *B. dorsalis* complex, involving not only *B. invadens* but also *B. dorsalis*, *B. papayae*, *B. philippinensis* and *B. carambolae*, all of which rank among the world's most destructive and highly invasive alien pest species. At the request from member states, the Joint FAO/IAEA Division took on an instrumental role in establishing in 2010 a six-year Coordinated Research Project (CRP) on '*Resolution of Cryptic Species Complexes of Tephritid Flies to Overcome Constraints to SIT Application and International Trade*'. While the first meeting of re-

searchers to coordinate the research programme of this CRP took place in Vienna in 2010, it must be noted that it was during the FAO/IAEA consultants meeting in 2009 (that included Jorge Hendrichs and Keng-Hong Tan) that the CRP was designed.

The objectives of the CRP were not only to resolve the species issues within the *B. dorsalis* complex but also of other tephritid genera where close species relationships were an issue, including *Anastrepha* and *Ceratitis* pest populations or species. Research co-ordination meetings were held in Vienna, Austria (2010), Brisbane, Australia, (2012), Tucuman, Argentina (2013), and La Reunion, France (2015), where research progress was critically evaluated and follow-up research work-plans developed. A multidisciplinary consensus was finally reached that while there was sufficient evidence confirming the separate species status for *B. carambolae*, the four species *B. dorsalis*, *B. invadens*, *B. papayae* and *B. philippinensis* constituted only a single biological species. This culminated in the synonymization of the later three species with *B. dorsalis* (Schutze et al. 2015a). The large authorship of the paper (49 authors from 15 countries) attests to the commitment, dedication and involvement of the international community to resolving this issue. As this outcome is celebrated, it must be remembered that doing good science through perseverance is not the only prerequisite for success, but strong scientific leadership in adopting the philosophy that asking and pursuing answers to the right questions is of paramount importance. Finally, we acknowledge that more research needs to be conducted to fill gaps in the biological, ecological and evolutionary understanding of other sibling species within the *B. dorsalis* complex, particularly incipient species in the Asia-Pacific region.

Acknowledgements

The authors would like to thank everyone in the *Bactrocera dorsalis* complex group in the CRP for the valuable discussions. We are also grateful to Alfred M. Handler (USDA, Gainesville, Florida) for his interest and invaluable assistance in conducting the *piggyBac* studies on *B. papayae*. The assistance of technical staff and interest of colleagues (Sai-Tee Tan) in the School of Biological Sciences, Universiti Sains Malaysia (1981–1999) are appreciated. Also thanks to Russell H. Messing (University of Hawaii) for providing the initial stock of the *B. dorsalis* s.s. to Keng-Hong Tan. The financial support from FAO/IAEA offered to Alvin K.W. Hee through IAEA research contract no. 16066 and Keng-Hong Tan, IAEA research contract no. 16160 is gratefully acknowledged.

References

- Armstrong KF, Cameron CM (2000) Species identification of tephritids across a broad taxonomic range using ribosomal DNA. In: Tan KH (Ed.) Area-wide Control of Fruit Flies and Other Insect Pests. Penang, Penerbit Universiti Sains Malaysia, Penang, 703–710.

- Behura SK (2006) Molecular marker systems in insects: current trends and future avenues. *Molecular Ecology* 15: 3087–3113. doi: 10.1111/j.1365-294X.2006.03014.x
- Cantrell BK, Chadwick B, Cahill A (2002) *Fruit Fly Fighters – Eradication of the Papaya Fruit Fly*, CSIRO Publishing, Collingwood, 208 pp.
- Clarke AR, Schutze MK (2014) The complexities of knowing what it is you are trapping. In: Shelly TE, Epsky N, Jang EB, Reyes-Flores J, Vargas RI (Eds) *Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies*. Springer, Dordrecht, 611–632. doi: 10.1007/978-94-017-9193-9_18
- Clarke AR, Allwood A, Chinajariyawong A, Drew RAI, Hengsawad C, Jirasurat M, Krong CK, Kritsaneepailoon S, Vijaysegaran S (2001) Seasonal abundance and host use patterns of seven *Bactrocera* Macquart species (Diptera: Tephritidae) In: Thailand and Peninsular Malaysia. *The Raffles Bulletin of Zoology* 49: 207–220.
- Clarke AR, Armstrong KF, Carmichael AE, Mine JR, Raghu S, Roderick GK, Yeates DK (2005) Invasive phytophagous pests arising through a recent tropical evolutionary radiation: the *Bactrocera dorsalis* complex of fruit flies. *Annual Review of Entomology* 50: 293–319. doi: 10.1146/annurev.ento.50.071803.130428
- Cruickshank L, Jessup AJ, Cruickshank DJ (2001) Interspecific crosses of *Bactrocera tryoni* (Froggatt) and *Bactrocera jarvisi* (Tryon) (Diptera: Tephritidae) in the laboratory. *Australian Journal of Entomology* 40: 278–280. doi: 10.1046/j.1440-6055.2001.00223.x
- Cunningham RT (1989) Parapheromones. In: Robinson AS, Hooper G (Eds) *World Crop Pests: Fruit flies. Their Biology, Natural Enemies and Control*, Vol. 3A, Elsevier, Amsterdam, 221–230.
- De Meyer M, Robertson MP, Mansell MW, Ekesi S, Tsuruta K, Mwaiko W, Vayssières JF, Peterson AT (2010) Ecological niche and potential geographic distribution of the invasive fruit fly *Bactrocera invadens* (Diptera, Tephritidae). *Bulletin of Entomological Research* 100: 35–48. doi: 10.1017/S0007485309006713
- Drew RAI (1989) The tropical fruit flies (Diptera: Tephritidae: Dacinae) of the Australasian and Oceanian regions. *Memoirs of the Queensland Museum* 26: 1–521.
- Drew RAI (1991) Taxonomic studies on oriental fruit fly. In: Vijaysegaran S, Ibrahim AG (Eds) *Fruit Flies in the Tropics*. Proceedings of the 1st International Symposium on Fruit Flies in the Tropics, Malaysian Agricultural Research and Development Institute and Malaysian Plant Protection Society, Kuala Lumpur, 63–66.
- Drew RAI (1997) The economic and social impact of the *Bactrocera papayae* Drew and Hancock (Asian papaya fruit fly) outbreak in Australia. In: Allwood AJ, Drew RAI (Eds) *Symposium on Management of Fruit Flies in the Pacific*. ACIAR Proceedings No. 76: 205–207.
- Drew RAI, Hancock DL (1994) The *Bactrocera dorsalis* complex of fruit flies (Diptera: Tephritidae: Dacinae) in Asia. *Bulletin of Entomological Research*, Supplement 2, 68 pp. doi: 10.1017/S1367426900000278
- Drew RAI, Romig MC (2013) *Tropical Fruit Flies of South-East Asia* (Tephritidae: Dacinae). CABI, Wallingford, 653 pp.
- Drew RAI, Tsuruta K, White IM (2005) A new species of pest fruit fly (Diptera: Tephritidae: Dacinae) from Sri Lanka and Africa. *African Entomology* 13: 149–154.

- Drew RAI, Raghu S, Halcoop P (2008) Bridging the morphological and biological species concepts: studies on the *Bactrocera dorsalis* (Hendel) complex (Diptera: Tephritidae: Dacinae) in South-east Asia. *Biological Journal of the Linnean Society* 93: 217–226. doi: 10.1111/j.1095-8312.2007.00952.x
- Fay HAC, Drew RAI, Lloyd AC (1997) The eradication program for papaya fruit fly (*Bactrocera papayae* Drew and Hancock) in North Queensland. In: Allwood AJ, Drew RAI (Eds) *Symposium on Management of Fruit Flies in the Pacific*. ACIAR Proceedings No. 76: 259–261.
- Handler AM, Zimowska GJ, Amstrong KF (2008) Highly similar piggyBac elements in *Bactrocera* that share a common lineage with elements in noctuid moths. *Insect Molecular Biology* 17: 387–393. doi: 10.1111/j.1365-2583.2008.00813.x
- Hardy DE, Adachi MS (1954) Studies in the fruit flies of the Philippines, Indonesia and Malaya Part 1. Dacini (Tephritidae-Diptera). *Pacific Science* 8: 147–204.
- He M, Haymer DS (1997) Polymorphic intron sequences detected within and between populations of the Oriental fruit fly (Diptera: Tephritidae). *Annals of the Entomological Society of America* 90: 825–831. doi: 10.1093/aesa/90.6.825
- He M, Haymer DS (2003) The use of intron sequences for molecular taxonomic and systematic studies of species in the *Bactrocera dorsalis* complex. *Plant Protection Bulletin* 5: 35–47.
- Hee AKW, Tan KH (1998) Attraction of female and male *Bactrocera papayae* to conspecific males fed with methyl eugenol and attraction of females to male sex pheromone components. *Journal of Chemical Ecology* 24: 753–764. doi: 10.1023/A:1022302605357
- Hee AKW, Tan KH (2004) Male sex pheromonal components derived from methyl eugenol in the hemolymph of the fruit fly *Bactrocera papayae*. *Journal of Chemical Ecology* 30: 2127–2138. doi: 10.1023/B:JOEC.0000048778.02561.70
- Hee AKW, Tan KH (2005) Bioactive fractions containing methyl eugenol-derived sex pheromonal components in haemolymph of the male fruit fly *Bactrocera dorsalis* (Diptera: Tephritidae). *Bulletin of Entomological Research* 95: 615–620. doi: 10.1079/BER2005392
- Hee AKW, Tan KH (2006) Transport of methyl eugenol-derived sex pheromonal components in the male fruit fly, *Bactrocera dorsalis*. *Comparative Biochemistry and Physiology, Part C: Toxicology and Pharmacology* 143: 422–428. doi: 10.1016/j.cbpc.2006.04.009
- Hill MP, Terblanche JS (2014) Niche overlap of congeneric invaders supports a single-species hypothesis and provides insight into future invasion risk: implications for global management of the *Bactrocera dorsalis* complex. *PLoS ONE* 9: e90121. doi: 10.1371/journal.pone.0090121
- Howlett FM (1915) Chemical reactions of fruit flies. *Bulletin of Entomological Research* 6: 297–305. doi: 10.1017/S0007485300043571
- Iwahashi O (2001) Aedeagal length of the Oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae), and its sympatric species in Thailand and the evolution of a longer and shorter aedeagus in the parapatric species of *B. dorsalis*. *Applied Entomology and Zoology* 36: 289–297. doi: 10.1303/aez.2001.289
- Kawasaki K, Iwahashi O, Kaneshiro KY (Eds) (1991) *Proceedings of the International Symposium on the Biology and Control of Fruit Flies, Okinawa (Japan), September 1991*. Food and Fertilizer Technology Center, 360 pp.

- Khamis FM, Masiga DK, Mohamed SA, Salifu D, De Meyer M, Ekesi S (2012) Taxonomic identity of the invasive fruit fly pest, *Bactrocera invadens*: concordance in morphometry and DNA barcoding. PLoS ONE 7: e44862. doi: 10.1371/journal.pone.0044862
- Khoo CCH, Yuen KH, Tan KH (2000) Attraction of female *Bactrocera papayae* to sex pheromone components with two different release devices. Journal of Chemical Ecology 26: 2487–2496. doi: 10.1023/A:1005576427138
- Li Y, Wu Y, Chen H, Wu J, Li Z (2012) Population structure and colonization of *Bactrocera dorsalis* (Diptera: Tephritidae) in China, inferred from mtDNA COI sequences. Journal of Applied Entomology 136: 241–251. doi: 10.1111/j.1439-0418.2011.01636.x
- Lux SA, Copeland RS, White IM, Manrakhan A, Billah M (2003) A new invasive fruit fly species from the *Bactrocera dorsalis* (Hendel) group detected in East Africa. Insect Science and its Application 23: 355–361. doi: 10.1017/S174275840001242X
- Mayr E (1957) Species concepts and definitions. In: Mayr E (Ed.) The species problem. American Association for Advancement of Science, Washington, DC, 1–22.
- McInnis DO, Rendon P, Jang EB, Van Saruers-Muller A, Sugayama R, Malavasi A (1999) Interspecific mating of introduced, sterile *Bactrocera dorsalis* with wild *B. carambolae* (Diptera: Tephritidae) in Suriname: a potential case for cross-species sterile insect technique. Annals of the Entomological Society of America 92: 758–65. doi: 10.1093/aesa/92.5.758
- Metcalfe RL, Mitchell WC, Fukuto TR, Metcalfe ER (1975) Attraction of the oriental fruit fly, *Dacus dorsalis*, to methyl eugenol and related olfactory stimulants. Proceedings of the National Academy of Sciences of the United States 72: 2501–2505. doi: 10.1073/pnas.72.7.2501
- Muraji M, Nakahara S (2001) Phylogenetic relationships among fruit flies, *Bactrocera* (Diptera, Tephritidae), based on the mitochondrial rDNA sequences. Insect Molecular Biology 10: 549–559. doi: 10.1046/j.0962-1075.2001.00294.x
- Naeole CKM, Haymer DS (2003) Use of oligonucleotide arrays for molecular taxonomic studies of closely related species in the oriental fruit fly (*Bactrocera dorsalis*) complex. Molecular Ecology Notes 3: 662–665. doi: 10.1046/j.1471-8286.2003.00545.x
- Nishida R, Tan KH, Fukami H (1988a) Cis-3,4-dimethoxycinnamyl alcohol from the rectal glands of male Oriental fruit fly, *Dacus dorsalis*. Chemistry Express 3: 207–210.
- Nishida R, Tan KH, Serit M, Lajis NH, Sukari AM, Takahashi S, Fukami H (1988b) Accumulation of phenylpropanoids in the rectal glands of males of the Oriental fruit fly, *Dacus dorsalis*. Experientia 44: 534–536. doi: 10.1007/BF01958941
- Schutze MK, Aketarawong N, Amornsak W, Armstrong KF, Augustinos A, Barr N, Bo W, Bourtzis K, Boykin LM, Cáceres C, Cameron SL, Chapman TA, Chinvinijkul S, Chomič A, De Meyer M, Drosopoulou ED, Englezou A, Ekesi S, Gariou-Papalexiou A, Hailstones D, Haymer D, Hee AKW, Hendrichs J, Hasanuzzaman M, Jessup A, Khamis FM, Krosch MN, Leblanc L, Mahmood K, Malacrida AR, Mavragani-Tsipidou P, McInnis DO, Mwatawala M, Nishida R, Ono H, Reyes J, Rubinoff DR, San Jose M, Shelly TE, Srikachar S, Tan KH, Thanaphum S, Ul Haq I, Vijaysegaran S, Wee SL, Yesmin F, Zacharopoulou A, Clarke AR (2015a) Synonymization of key pest species within the *Bactrocera dorsalis* complex (Diptera: Tephritidae): taxonomic changes based on 20 years of integra-

- tive morphological, genetic, behavioural, and chemoecological data. *Systematic Entomology* 40: 456–471. doi: 10.1111/syen.12113
- Schutze MK, Mahmood K, Pavasovic A, Bo W, Newman J, Clarke AR, Krosch MN, Cameron SL (2015b) One and the same: integrative taxonomic evidence that the African Invasive Fruit Fly *Bactrocera invadens* (Diptera: Tephritidae), is the same species as the Oriental Fruit Fly *Bactrocera dorsalis*. *Systematic Entomology* 40: 472–486. doi : 10.1111/syen.12114
- Shelly TE, Epsky N, Jang EB, Reyes-Flores J, Vargas RI (Eds) (2014) *Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies*, Springer, Dordrecht, 638 pp. doi: 10.1007/978-94-017-9193-9
- Smith PT, Kambhampati S, Armstrong KF (2003) Phylogenetic relationships among *Bactrocera* species (Diptera: Tephritidae) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 26: 8–7. doi: 10.1016/S1055-7903(02)00293-2
- Tan KH (1984) Description of a new attractant trap and the effect of placement height on two *Dacus* species (Diptera: Tephritidae). *Journal of Plant Protection in the Tropics* 1: 117–120.
- Tan KH (1985) Estimation of native populations of male *Dacus* spp. by using Jolly's stochastic method using a new designed attractant trap in a village ecosystem. *Journal of Plant Protection in the Tropics* 2: 87–95.
- Tan KH (2000) Behaviour and chemical ecology of *Bactrocera* flies. In: Tan KH (Ed.) *Area-Wide Control of Fruit Flies and Other Insect Pests*. Penerbit Universiti Sains Malaysia, Penang, 647–656.
- Tan KH (2003) Interbreeding and DNA analysis of sibling species within the *Bactrocera dorsalis* complex. In: *Recent Trends on Sterile Insect Technique and Area-Wide Integrated Pest Management- Economic Feasibility, Control Projects, Farmer Organization and Bactrocera dorsalis Complex Control Study*. Research Institute for Subtropics, Okinawa, 113–122.
- Tan KH, Lee SL (1982) Species diversity and abundance of *Dacus* (Diptera: Tephritidae) in five ecosystems of Penang, West Malaysia. *Bulletin of Entomological Research* 72: 709–716. doi: 10.1017/S0007485300008737
- Tan KH, Jaal Z (1986) Comparison of male adult population densities of the Oriental and Artocarpus fruit flies *Dacus* spp. (Diptera: Tephritidae), in two nearby villages in Penang, Malaysia. *Researches in Population Ecology* 28: 85–89. doi: 10.1007/BF02515538
- Tan KH, Serit M (1988) Movement and population density comparison of native male adult *Dacus dorsalis* and *Dacus umbrosus* (Diptera: Tephritidae) among three ecosystems. *Journal of Plant Protection in the Tropics* 5: 17–21.
- Tan KH, Nishida R (1996) Sex pheromone and mating competition after methyl eugenol consumption in the *Bactrocera dorsalis* complex. In: McPherson BA, Steck GJ (Eds) *Fruit Fly Pests. A World Assessment of Their Biology and Management*. St. Lucie Press, Florida, 147–153.
- Tan KH, Nishida R (2012) Methyl eugenol- its occurrence, distribution, and role in nature, especially in relation to insect behavior and pollination. *Journal of Insect Science* 20: 74 pp. doi: 10.1673/031.012.5601
- Tan KH, Kirton LG, Serit M (1987) Age response of *Dacus dorsalis* (Hendel) to methyl eugenol in (a) a wind tunnel, and (b) traps set in a village, and its implication in population

- estimation. In: Economopoulos AP (Ed.) Fruit Flies. Proceedings of the 2nd International Symposium, Colymbari (Greece), September 1986. G. Tsiveriotis Ltd, Athens, 425–432.
- Tan KH, Nishida R, Jang EB, Shelly TE (2014) Pheromones, male lures, and trapping of tephritid fruit flies. In: Shelly TE, Epsky N, Jang EB, Flores-Reyes J, Vargas RI (Eds) Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies, Springer, New York, 15–74. doi: 10.1007/978-94-017-9193-9_2
- Vargas RI, Shelly TE, Leblanc L, Piñero JC (2010) Recent advances in methyl eugenol and cue-lure technologies for fruit fly detection, monitoring, and control in Hawaii. In: Litwack G (Ed.) Vitamins and Hormones. Volume 83, Academic, Burlington, 575–596. doi: 10.1016/S0083-6729(10)83023-7
- Verghese A, Tandon PL, Stonehouse JM (2004) Economic evaluation of the integrated management of the oriental fruit fly *Bactrocera dorsalis* (Diptera: Tephritidae) in Mango in India. Crop Protection 23: 61–63. doi: 10.1016/S0261-2194(03)00087-5
- Wee SL (2000) Behaviour and reproductive ecology of fruit fly, *Bactrocera carambolae* and *B. papayae*. PhD thesis, Universiti Sains Malaysia, Penang. [In Malay]
- Wee SL, Tan KH (2000) Interspecific mating of two sibling species of the *Bactrocera dorsalis* complex in a field cage. In: Tan KH (Ed.) Area-Wide Control of Fruit Flies and Other Insect Pests, Penerbit Universiti Sains Malaysia, Penang, 667–674.
- Wee SL, Tan KH (2005) Evidence of natural hybridization between two sympatric sibling species of *Bactrocera dorsalis* complex based on pheromone analysis. Journal of Chemical Ecology 31: 845–858. doi 10.1007/s10886-005-3548-6
- Wee SL, Tan KH (2007) Temporal accumulation of phenylpropanoids in male fruit flies, *Bactrocera dorsalis* and *B. carambolae* (Diptera: Tephritidae) following methyl eugenol consumption. Chemoecology 17: 81–85. doi: 10.1007/s00049-006-0363-7
- White IM, Elson-Harris MM (1992) Fruit flies of economic significance: Their identification and bionomics. CAB International, Oxford, 601 pp.
- Ye H (2001) Distribution of the oriental fruit fly (Diptera: Tephritidae) in Yunnan Province. Insect Science 8: 175–182. doi: 10.1111/j.1744-7917.2001.tb00483.x
- Yong HS (1994) Hydroxybutyrate dehydrogenase electromorph distinguishes *Bactrocera dorsalis* (Insecta: Diptera: Tephritidae) from sibling taxa in Peninsular Malaysia. In: Yong HS, Khoo SG (Eds) Current research on tropical fruit flies and their management, The Working Group on Malaysian Fruit Flies, Kuala Lumpur, 23–26.
- Yong HS (1995) Genetic differentiation and relationship in five taxa of the *Bactrocera dorsalis* complex (Insecta: Diptera: Tephritidae). Bulletin of Entomological Research 85: 431–435. doi: 10.1017/S0007485300036166
- Zimowska GJ, Handler AM (2005) *PiggyBac* transposons in *Bactrocera dorsalis* and their use for species identification. 8th Exotic Fruit Fly Symposium. Riverside, California, March 7–9, 2005.